


ADVANTAGE FOR CATS



For use on cats only. Do not use on dogs or rabbits.
For use on kittens and cats from 8 weeks of age.

INDICATION

- A fast action, monthly spot-on flea treatment.
- Kills 98-100% adult fleas within half a day of administration.
- Protects the cats by killing and controlling fleas.
- Starts working within half an hour of application.
- Stops flea biting within 3-5 minutes.
- Prevents diseases like Flea Allergy Dermatitis (FAD).
- Waterproof treatment that remains effective even after bathing or swimming.

- Safe to be administered on nursing cats and queens.
 - Contains Imidacloprid as an active ingredient that kills adult fleas very swiftly thus freeing the cat from the irritation caused by it.
 - Very easy to administer on cats.
-

COMPOSITION

Imidacloprid	10% w/v
--------------	---------

RESIDUAL EFFECT

A monthly spot-on flea preventive treatment that stops flea bites within 3 to 5 minutes of application thereby killing adult fleas and larvae within 20 minutes of administration. It also reduces the outbreak of Flea Allergy Dermatitis (FAD).

WARNING

- Not intended for human use. Keep it away from children.
 - Not to be used for any purpose, or in any manner, contrary to this unless authorized under appropriate legislation.
-

PRECAUTIONS

- If your cat is already suffering from a disease do not administer this treatment on her.
- Never administer this treatment on dogs.

- Check the expiry date of the product before administering it.
 - Children and other pets should not be allowed to get near the treated pet.
-

DOSAGE

CAT WEIGHT	MONTHLY DOSE
1 to 10 lbs. (S)	0.4 ml tube of imidacloprid 100g/L
more than 10 lbs.	0.8 ml tube of imidacloprid 100g/L

METHOD OF APPLICATION

- Use as directed by your veterinarian.
- Recommended for kittens and cats at least 8 weeks of age.
- Hold the applicator tube in an erect position. Unlock the tube cap.
- Turn the cap around and place the other end of the cap back on the tube.
- Twirl the cap and open the seal then get rid of the cap from the tube.



Step 1

Keep the kitty in an erect position.



Step 2

Part her hair on the back of the neck at the base of the skull so that you can see the skin completely. Keep the tip of the tube on the skin and squeeze out the content completely to smear the solution directly on the skin.

- Let the solution dry out completely before allowing children and other pets to touch the applied spot.
-

STORAGE AND DISPOSAL

- **Storage:** Store at a room temperature below 30 degrees Celsius.
 - **Disposal:** Dispose of used tubes and packaging by wrapping in paper and placing in garbage.
-

PESTICIDE DISPOSAL

- Do not contaminate water, food, or feed by storage and disposal.
 - Never empty product in indoor or outdoor drain.
-

CONTAINER DISPOSAL

- **If the container is empty:** Never re-use or refill the container. Keep it in trash or offer for recycling if the option is available.
 - **If the container is wholly filled:** Contact your local solid waste agency for disposal instructions. Never keep the new product near any indoor or outdoor drain.
-

FIRST AID

Keep the product container or label with you while calling a poison control center or doctor or while going for the treatment:

- **If the treatment is swallowed:** Contact poison contact center or doctor quickly for getting their advice. If the person is able to swallow, give him/her a sip of glass of water. Never induce vomiting until and unless it has been recommended by the poison center or doctor. If the person is unconscious do not give anything by mouth.
- **If the treatment gets into the eyes:** Keep the eye open and rinse slowly and softly with water for 15 to 20 minutes. If the person is wearing contact lenses, remove it after 5 minutes and continue rinsing the eye. Call the doctor or poison control center for advice.
- **If the treatment affects the skin:** Wash the spot with plenty of soap and water.
- **For the physician:** Treat the patient systematically.

SIDE EFFECTS

Although Advantage Flea treatment is considered to be safe, certain cats may develop an adverse reaction to the treatment. Some of the side effects of this monthly spot-on treatment are:

- Falling of hair out of the cat on the application site (***But this is not considered to be a major issue since the hair usually grow back uneventfully.***)

- If the cat licks the application site she will often salivate profusely with distaste, (***cats rarely consume enough of imidacloprid insecticide to become poisoned***) but if the owner notices that a large quantity of the product has been consumed by the kitty, immediate veterinary attention would be required.

Manufactured by

Bayer



Visit our website at -

<https://www.canadavetexpress.com/flea-and-tick-control-treatment/advantage-for-cats/p23.aspx>

DISCLAIMER

The information provided in this file is for informational purpose only. Please consult your veterinarian before administering the product to your pet.

©2016 CanadaVetExpress.com - All Rights Reserved