



Canada 
VetExpress.com
Care - Quality - Service

ADVANTAGE FOR DOGS



For use on dogs only. Do not use on cats or rabbits.

For use on puppies from 8 weeks of age

INDICATION

- A spot-on insecticide for flea control in dogs and cats.
- Indicated for use in dogs at least 8 weeks of age.
- Extremely safe for puppies, lactating dogs along with pregnant dogs.
- Indicated for the prevention of fleas thereby controlling the flea population.
- Assists in treating Flea Allergy Dermatitis (FAD) in dogs.
- Waterproof property enables pets to easily go in water without worrying about the treatment to cease its effect.
- Contains Imidacloprid that stops flea bites within 3 to 5 minutes of application thereby killing adult fleas and larvae within 20 minutes of administration.

- Does not require fleas to bite the pet to produce the desired results.
 - Assists in reducing the outbreak of Flea Allergy Dermatitis (FAD).
-

COMPOSITION

Imidacloprid	10% w/v
--------------	---------

RESIDUAL EFFECT

A monthly spot-on flea preventive treatment that stops flea bites within 3 to 5 minutes of application thereby killing adult fleas and larvae within 20 minutes of administration. It also reduces the outbreak of Flea Allergy Dermatitis (FAD).

WARNING

Not intended for human use. Keep it away from children.

PRECAUTIONS

The safety of this product on debilitated or aged dog has not been verified yet. So, consult with your veterinarian before administering this treatment on him/her.

DOSAGE

DOG WEIGHT	MONTHLY DOSE
1 to 10 lbs.	0.4 ml tube
11 to 20 lbs.	1.0 ml tube
21 to 55 lbs.	2.5 ml tube
over 55 lbs.	4 ml tube

METHOD OF APPLICATION

- Use as directed by your veterinarian.
- Recommended for puppies and dogs at least 8 weeks of age.
- Hold the applicator tube in an erect position. Unlock the tube cap.
- Turn the cap around and place the other end of the cap back on the tube.
- Twirl the cap and open the seal then get rid of the cap from the tube.



Step 1

Keep the doggy in an erect position.



Step 2

Part his hair on the back of the neck at the base of the dog's skull so that you can see the skin completely. Keep the tip of the tube on the skin and squeeze out the content completely to smear the solution directly on the dog's skin.

- Let the solution dry out completely before allowing children and other pets to touch the applied spot.
-

STORAGE AND DISPOSAL

Never pollute water, food or feed through storage or disposal.

PESTICIDE DISPOSAL

- Do not contaminate water, food, or feed by storage and disposal.
 - Never empty product in indoor or outdoor drain.
-

CONTAINER DISPOSAL

- **If the container is empty:** Never re-use or refill the container. Keep it in trash or offer for recycling if the option is available.
 - **If the container is wholly filled:** Contact your local solid waste agency for disposal instructions. Never keep the new product near any indoor or outdoor drain.
-

FIRST AID

Keep the product container or label with you while calling a poison control center or doctor or while going for the treatment:

- **If the treatment is swallowed:** Contact poison contact center or doctor quickly for getting their advice. If the person is able to swallow, give him/her a sip of glass of water. Never induce vomiting until and unless it has been recommended by the poison center or doctor. If the person is unconscious do not give anything by mouth.

- **If the treatment gets into the eyes:** Keep the eye open and rinse slowly and softly with water for 15 to 20 minutes. If the person is wearing contact lenses, remove it after 5 minutes and continue rinsing the eye. Call the doctor or poison control center for advice.

 - **If the treatment affects the skin:** Wash the spot with plenty of soap and water.

 - **For the physician:** Treat the patient systematically.
-

SIDE EFFECTS

Although Advantage Flea treatment is considered to be safe, certain dogs may develop an adverse reaction to the treatment. Since it is applied on the skin of the doggy, it gets absorbed into the skin and enters his body. It can then get into the circulatory system of the pet and cause brain damage and nerve damage.

Watch out for these symptoms:

- Biting
- Chewing at the application site
- Diarrhea
- Difficulty in breathing
- Dilation of the pupils
- Increased thirst
- Itching
- Excessive salivation or drooling
- Irritation of the eyes
- Lethargy
- Loss of appetite

- Non-responsiveness
- Rash at the application site
- Seizures
- Shaking uncontrollably
- Skin irritation
- Vomiting

Manufactured by
Bayer



Visit our website at

<https://www.canadavetexpress.com/flea-and-tick-control-treatment/advantage-for-dogs/p24.aspx>

DISCLAIMER

The information provided in this file is for informational purpose only. Please consult your veterinarian before administering the product to your pet.

©2016 CanadaVetExpress.com - All Rights Reserved